ELSEVIER

Environmental Pollution 147 (2007) I–VII

ENVIRONMENTAL POLLUTION

ww.elsevier.com/locate/envpol

Contents of Volume 147

Number 1 2007

Commentary

1 **Where do Chinese scientists publish their research in environmental science and technology?**
Y.-G. Zhu, P. O'connor, J.-H. Cao

Regular Papers

4 **The influence of maize residues on the mobility and binding of benazolin: Investigating physically extracted soil fractions**
F. Schnitzler, A. Lavorenti, A.E. Berns, N. Drewes, H. Vereecken, P. Burauel

Addition of crop residues increases the amount of non-extractable residues and decreases the mobility of benazolin and its metabolites.

14 **Collembola and macroarthropod community responses to carbamate, organophosphate and synthetic pyrethroid insecticides: Direct and indirect effects**
G.K. Frampton, P.J. van den Brink

Direct and indirect insecticide effects differ among closely-related arthropod taxa; resurgence of Collembola may occur widely after synthetic pyrethroid insecticide applications.

26 **Field effects of pollutants in dynamic environments. A case study on earthworm populations in river floodplains contaminated with heavy metals**
C. Klok, P.W. Goedhart, B. Vandecasteele

Interaction with flooding masks heavy metal effects.

32 **Distribution and movement of nutrients and metals in a *Pinus radiata* forest soil following applications of biosolids**
R.G. McLaren, L.M. Clucas, T.W. Speir, A.P. van Schaik

Biosolids application to forest soils results in substantial build-up of macronutrients and metals in the forest litter layer.

41 **Study on adsorption behavior of volatile and semivolatile organic vapors to air-dry soils based on QSPR methods**
H. Liu, X. Yao, M. Liu, Z. Hu, B. Fan

The model we built can be used to predict a given molecule's adsorption characteristics, and also allow a more accurate determination of its environmental fate.

50 **Localisation and quantification of elements within seeds of Cd/Zn hyperaccumulator *Thlaspi praecox* by micro-PIXE**
K. Vogel-Mikuš, P. Pongrac, P. Kump, M. Nečemer, J. Simčič, P. Pelicon, M. Budnar, B. Povh, M. Regvar

Hyperaccumulation of Cd with partitioning in the epidermis of cotyledons was observed in T. praecox embryos.

60 **Diclofenac disposition in Indian cow and goat with reference to *Gyps* vulture population declines**
M.A. Taggart, R. Cuthbert, D. Das, C. Sashikumar, D.J. Pain, R.E. Green, Y. Feltrer, S. Shultz, A.A. Cunningham, A.A. Meharg

Residual diclofenac in livestock carcasses in India poses a continuing risk to Gyps vultures.

66 **Efficiency of surfactant-enhanced desorption for contaminated soils depending on the component characteristics of soil-surfactant–PAHs system**
W. Zhou, L. Zhu

The efficiency of surfactant-enhanced desorption showed strong dependence on the soil composition, surfactant structure and compounds properties.

74 **Phytoremediation of mixed-contaminated soil using the hyperaccumulator plant *Alyssum lesbiacum*: Evidence of histidine as a measure of phytoextractable nickel**
A.C. Singer, T. Bell, C.A. Heywood, J.A.C. Smith, I.P. Thompson

Alyssum lesbiacum was shown to phytoextract nickel from PAH-contaminated soils from which the pool of nickel accessed for phytoextraction is closely modelled by a histidine-soil extract.

83 **Effect of temperature and season on reproduction, neutral red retention and metallothionein responses of earthworms exposed to metals in field soils**
C. Svendsen, P.K. Hankard, L.J. Lister, S.K. Fishwick, M.J. Jonker, D.J. Spurgeon

Laboratory and field studies demonstrate metal effects on earthworm life-cycle and biochemical responses are not influenced by temperature regime.

94 **The antioxidative response system in *Glycine max* (L.) Merr. exposed to Deltamethrin, a synthetic pyrethroid insecticide**
F. Bashir, Mahmooduzzafar, T.O. Siddiqi, M. Iqbal

The Deltamethrin-induced oxidative stress alters the ascorbate–glutathione cycle in Glycine max.

doi:10.1016/S0269-7491(07) 00171-6

101 Spatial relationships between nitrogen status and pitch canker disease in slash pine planted adjacent to a poultry operation
I. Lopez-Zamora, C. Bliss, E.J. Jokela, N.B. Comerford, S. Grunwald, E. Barnard, G.M. Vasquez

Local emissions from poultry production appear to significantly contribute to the spatial distribution of N and pitch canker disease in managed slash pine ecosystems.

112 Remediation of cadmium-contaminated paddy soils by washing with calcium chloride: Verification of on-site washing
T. Makino, T. Kamiya, H. Takano, T. Itou, N. Sekiya, K. Sasaki, Y. Maejima, K. Sugahara

In situ soil washing in a paddy field using an on-site wastewater treatment system resulted in an effective decrease of Cd in soil and rice grains without affecting rice yield.

120 Reconstructing temporal trends in heavy metal deposition: Assessing the value of herbarium moss samples
L. Shotbolt, P. Büker, M.R. Ashmore

Herbarium moss samples can contribute to the reconstruction of past heavy metal deposition.

131 Nutrient losses from manure and fertilizer applications as impacted by time to first runoff event
D.R. Smith, P.R. Owens, A.B. Leytem, E.A. Warnemuende

Fertilization with manures results in lower nutrient runoff than inorganic fertilizers, especially if at least one week passes between fertilization and runoff.

138 Reducing fluxes of faecal indicator compliance parameters to bathing waters from diffuse agricultural sources: The Brighouse Bay study, Scotland
D. Kay, M. Aitken, J. Crowther, I. Dickson, A.C. Edwards, C. Francis, M. Hopkins, W. Jeffrey, C. Kay, A.T. McDonald, D. McDonald, C.M. Stapleton, J. Watkins, J. Wilkinson, M.D. Wyer

Diffuse microbiological pollution from farming activities can be reduced by protected riparian zones.

150 *In situ* studies on the distribution patterns and dynamics of microcystins in a biomanipulation fish – bighead carp (*Aristichthys nobilis*)
J. Chen, P. Xie, D. Zhang, H. Lei

Bighead carp is resistant to microcystins and can be used as biomanipulation fish to counteract cyanotoxin contamination.

158 Mortality and toxin bioaccumulation in *Bufo marinus* following exposure to *Cylindrospermopsis raciborskii* cell extracts and live cultures
S.H. White, L.J. Duivenvoorden, L.D. Fabbro, G.K. Eaglesham

Cane toad tadpoles experienced adverse effects and bioaccumulation of toxin following exposure to the cyanobacterium Cylindrospermopsis raciborskii.

168 Heavy metal concentrations in soils and plant accumulation in a restored manganese mineland in Guangxi, South China
M.S. Li, Y.P. Luo, Z.Y. Su

Soils of a restored manganese mineland are basically unpolluted, but direct cultivation of edible agronomic crops may be a health risk for humans.

176 Non-target screening of extractable and non-extractable organic xenobiotics in riverine sediments of Ems and Mulde Rivers, Germany
A. Kronimus, J. Schwarzbauer

Analysis of both, extractable and non-extractable fraction of organic xenobiotics in sediments reveals a more detailed and comprehensive anthropogenic load profile.

187 Uptake of oxytetracycline and its phytotoxicity to alfalfa (*Medicago sativa* L.)
W.D. Kong, Y.G. Zhu, Y.C. Liang, J. Zhang, F.A. Smith, M. Yang

Plant uptake of antibiotic oxytetracycline is energy-dependent.

194 What difference might sewage treatment performance make to endocrine disruption in rivers?
A.C. Johnson, R.J. Williams, P. Simpson, R. Kanda

Biological (trickling) filter sewage plants are less successful than other secondary sewage treatment plants at removing estrone from sewage effluent.

203 Characterization of TSP-bound *n*-alkanes and polycyclic aromatic hydrocarbons at rural and urban sites of Tianjin, China
S.-P. Wu, S. Tao, Z.-H. Zhang, T. Lan, Q. Zuo

Precipitation and sampling height can significantly influence TSP and TSP-bound PAHs.

211 Atmospheric concentration characteristics and gas–particle partitioning of PCBs in a rural area of eastern Germany
M. Mandalakis, E.G. Stephanou

Volatilization from terrestrial surfaces contaminated with Clophen A30 is still the main source that controls the atmospheric levels of PCBs in rural eastern Germany.

222 Differential responses of the freshwater wetland species *Juncus effusus* L. and *Caltha palustris* L. to iron supply in sulfidic environments
M.E.W. Van der Welle, K. Niggebrugge, L.P.M. Lamers, J.G.M. Roelofs

Toxicity of iron and sulfide are interacting with each other and have the potential to affect vegetation composition.

231 Heavy metal phytoremediation from a meta-analytical perspective
P. Audet, C. Charest

This meta-analytical approach has revealed a compromise between growth and HM uptake when plants are subjected to toxic soil-HM levels.

238 Magnetic approach to normalizing heavy metal concentrations for particle size effects in intertidal sediments in the Yangtze Estuary, China
W. Zhang, L. Yu, M. Lu, S.M. Hutchinson, H. Feng

Heavy metal concentration normalization for particle size effects by a magnetic approach facilitating sediment-based environmental monitoring.

245 Heavy metal load and dominance hierarchy in juvenile willow tits during winter
O. Hogstad, H.C. Pedersen

Cadmium concentrations in juvenile male willow tits (Parus montanus) were correlated with plasma testosterone levels.

248 Role of microbial inoculation and chitosan in phytoextraction of Cu, Zn, Pb and Cd by *Elsholtzia splendens* – a field case
F.Y. Wang, X.G. Lin, R. Yin

Co-application of microbial inocula and chitosan enhanced heavy metal phytoextraction by E. splendens.

256 Adsorption and absorption of dichlorodiphenyltrichloroethane (DDT) and metabolites (DDD and DDE) by rice roots
S.H. Chen, F.L. Xu, R. Dawson, X.C. Jiao, S. Tao

The adsorption and absorption of DDT by rice roots was quantitatively described by a three-step process model.

262 Chronic toxicity of arsenic, cobalt, chromium and manganese to *Hyalella azteca* in relation to exposure and bioaccumulation
W.P. Norwood, U. Borgmann, D.G. Dixon

Chronic toxicity of As, Co, Cr and Mn to Hyalella azteca can be described using a saturation-based mortality model in relationship to total-body or water metal concentration.

273 Active carbon-pools in rhizosphere of wheat (*Triticum aestivum* L.) grown under elevated atmospheric carbon dioxide concentration in a Typic Haplustept in sub-tropical India
P.C.B. Kant, S. Bhadraray, T.J. Purakayastha, V. Jain, M. Pal, S.C. Datta

The elevation in atmospheric CO_2 concentration can potentially increase the active carbon-pools in wheat rhizosphere in the semi-arid India.

282 The tidal hydrodynamics modeling of the Topolobampo coastal lagoon system and the implications for pollutant dispersion
Y. Montaño-Ley, R. Peraza-Vizcarra, F. Páez-Osuna

A model has been developed to simulate the tidal hydrodynamics and the behavior of a pollutant in the Topolobampo lagoon.

291 Accumulation of heavy metals in the Oostriku peat bog, Estonia: Determination of binding processes by means of sequential leaching
K. Syrovetnik, M.E. Malmström, I. Neretnieks

Sorption and precipitation processes are important in metal retention over a long time.

Number 2 2007

SPECIAL ISSUE SECTION: Environmental Pollution in China

Guest Editorial

301 China steps up its efforts in research and development to combat environmental pollution
Y.-G. Zhu, L. Wang, Z.-J. Wang, P. Christie, J.N.B. Bell

303 Source apportionment of polycyclic aromatic hydrocarbons in surface soil in Tianjin, China
Q. Zuo, Y.H. Duan, Y. Yang, X.J. Wang, S. Tao

PAHs in surface soil of Tianjin were apportioned and coal combustion, vehicle exhaust, coke production, and biomass burning were found to be the major sources.

311 Trace metal distribution in sediments of the Pearl River Estuary and the surrounding coastal area, South China
C.C.M. Ip, X.-D. Li, G. Zhang, O.W.H. Wai, Y.-S. Li

The distribution of trace metals in sediments reflected contaminant sources, physical and chemical deposition processes.

324 Response of *Microcystis* to copper stress – Do phenotypes of *Microcystis* make a difference in stress tolerance?
Z.-x. Wu, N.-q. Gan, Q. Huang, L.-r. Song

Stress-resistance ability in Microcystis is influenced by its phenotypes.

331 Occurrence of estrogenic effects in sewage and industrial wastewaters in Beijing, China
M. Ma, K. Rao, Z. Wang

Estrogenic activity in Beijing river water is attributed to direct discharges of untreated municipal and industrial wastewaters.

337 A biomimetic absorbent for removal of trace level persistent organic pollutants from water
H. Liu, J. Qu, R. Dai, J. Ru, Z. Wang

Triolein-embedded absorbent was developed and it could remove lipophilic pollutants from water effectively.

343 Generalized models for prediction of pentachlorophenol dissipation dynamics in soils
Y. He, J. Xu, H. Wang, Y. Wu

The empirical models were obtained for predicting PCP dissipation in soil.

350 Enhanced desorption of phenanthrene from contaminated soil using anionic/nonionic mixed surfactant
W. Zhou, L. Zhu

Anionic/nonionic mixed surfactants appear to be a better choice than single surfactants for the application of surfactant remediation technology.

358 Distribution of polycyclic aromatic hydrocarbons in thirty typical soil profiles in the Yangtze River Delta region, east China
L.F. Ping, Y.M. Luo, H.B. Zhang, Q.B. Li, L.H. Wu

The information was provided on levels, distributions, and possible sources of PAHs in the soils of Yangtze River Delta area.

366 **Adsorption and redox reactions of heavy metals on synthesized Mn oxide minerals**
X.H. Feng, L.M. Zhai, W.F. Tan, F. Liu, J.Z. He

The characteristics of heavy metal adsorption and Cr(III) oxidation on Mn oxide minerals are determined by their structure, composition, surface property and crystallinity.

374 **Effects of the arbuscular mycorrhizal fungus *Glomus mosseae* on growth and metal uptake by four plant species in copper mine tailings**
B.D. Chen, Y.-G. Zhu, J. Duan, X.Y. Xiao, S.E. Smith

This study demonstrated that AM associations can encourage plant survival in Cu mine tailings.

381 **Zinc and cadmium accumulation and tolerance in populations of *Sedum alfredii***
D.M. Deng, W.S. Shu, J. Zhang, H.L. Zou, Z. Lin, Z.H. Ye, M.H. Wong

Marked variation of Zn and Cd tolerance and accumulation exist in populations of Sedum alfredii, a Zn/Cd hyperaccumulator.

387 **Sulfur (S)-induced enhancement of iron plaque formation in the rhizosphere reduces arsenic accumulation in rice (*Oryza sativa* L.) seedlings**
Z.-Y. Hu, Y.-G. Zhu, M. Li, L.-G. Zhang, Z.-H. Cao, F.A. Smith

Sulfur-induced enhancement of iron plaque formation on the root surface of rice.

394 **Ground-level ozone in China: Distribution and effects on crop yields**
X. Wang, W. Manning, Z. Feng, Y. Zhu

More research is needed to reliably estimate ozone caused crop losses in China.

401 **Air quality impacts of power plant emissions in Beijing**
J. Hao, L. Wang, M. Shen, L. Li, J. Hu

CALMET/CALPUFF modeling estimates sizeable reductions in SO_2, PM_{10} and NO_x pollution from power plants by 2008 for Beijing.

409 **Photoelectrocatalytic degradation of pentachlorophenol in aqueous solution using a TiO_2 nanotube film electrode**
X. Quan, X. Ruan, H. Zhao, S. Chen, Y. Zhao

The characters and application influencing factors of the TiO_2 nanotube film electrode fabricated by the anodic oxidation method were investigated.

415 **Catalytic performance of Ag/Al_2O_3-C_2H_5OH-Cu/Al_2O_3 system for the removal of NOx from diesel engine exhaust**
C. Zhang, H. He, S. Shuai, J. Wang

Combination of Ag/Al_2O_3-C_2H_5OH-Cu/Al_2O_3 is a promising system for the removal of NO_x from diesel engine exhaust.

422 **Mechanisms of silicon-mediated alleviation of abiotic stresses in higher plants: A review**
Y. Liang, W. Sun, Y.-G. Zhu, P. Christie

This review article overviews roles Si plays in alleviating abiotic stress in higher plants and discusses future research directions.

Number 3 2007

SPECIAL ISSUE SECTION: Air Pollution and Climate Change: A Global Overview of the Effects on Forest Vegetation

Obituary

iii **Steve Hopkin 1956–2006**
M.T. Fountain, D. Spurgeon, C. Svendsen, H. Noel, N. Sorokin

429 **Introductory remarks to the special issue – XXII IUFRO World Congress, 2005 Brisbane – Air pollution and climate change: A global overview of the effects on forest vegetation**
M. Schaub, E. Paoletti

430 **Photosynthetic behavior of woody species under high ozone exposure probed with the JIP-test: A review**
F. Bussotti, R.J. Strasser, M. Schaub

Analytical techniques based on the direct fluorescence of chlorophyll a, allow us to discriminate species-specific physiological behavior in relation to ozone air pollution.

438 **Integrated effects of air pollution and climate change on forests: A northern hemisphere perspective**
A. Bytnerowicz, K. Omasa, E. Paoletti

Simultaneous addressing air pollution and climate change effects on forests is an opportunity for capturing synergies in future research and monitoring.

446 **A review of ozone-induced effects on the forests of central Mexico**
M.de L. de Bauer, T. Hernández-Tejeda

Ozone induced symptoms, poor tree regeneration and limited root colonization by mycorrhiza fungi observed in the valley of Mexico.

454 **Assessing the risk caused by ground level ozone to European forest trees: A case study in pine, beech and oak across different climate regions**
L.D. Emberson, P. Büker, M.R. Ashmore

A new flux-based model provides a revised assessment of risks of ozone impacts to European forests.

467 **Linking stress with macroscopic and microscopic leaf response in trees: New diagnostic perspectives**
M.S. Günthardt-Goerg, P. Vollenweider

Macroscopic leaf symptoms and their microscopic analysis as stress bioindications.

489 **Perspectives regarding 50 years of research on effects of tropospheric ozone air pollution on US forests**
D.F. Karnosky, J.M. Skelly, K.E. Percy, A.H. Chappelka

Elevated tropospheric ozone remains an important phytotoxic air pollutant over large areas of US forests.

507 **Impact of elevated CO_2 and nitrogen fertilization on foliar elemental composition in a short rotation poplar plantation**
S. Marinari, C. Calfapietra, P. De Angelis, G. Scarascia-Mugnozza, S. Grego

CO_2 enrichment reduced foliar nitrogen and increased the magnesium concentration in poplar.

516 **Photosynthetic responses to elevated CO_2 and O_3 in *Quercus ilex* leaves at a natural CO_2 spring**
E. Paoletti, G. Seufert, G. Della Rocca, H. Thomsen

Very high CO_2 concentrations did not compensate for the effects of O_3 on holm oak photosynthesis.

525 **Defense and avoidance of ozone under global change**
M. Tausz, N.E. Grulke, G. Wieser

Modeling of ozone effects on plants should include a measure for the plant defense capacity.

New Initiative

532 **Harmful effects of atmospheric nitrous acid on the physiological status of Scots pine trees**
H. Sakugawa, J.N. Cape

Exposure to HONO affects photosynthesis and nutrient status of pine trees.

Regular Papers

535 **Comparative toxicity of chlorpyrifos, diazinon, malathion and their oxon derivatives to larval *Rana boylii***
D.W. Sparling, G. Fellers

Laboratory tests on the toxicity of OP insecticides and their oxons suggest that they may be acutely lethal to amphibians at ecologically relevant concentrations.

540 **Tolerance of transgenic canola plants (*Brassica napus*) amended with plant growth-promoting bacteria to flooding stress at a metal-contaminated field site**
A.J. Farwell, S. Vesely, V. Nero, H. Rodriguez, K. McCormack, S. Shah, D.G. Dixon, B.R. Glick

Using transgenic plants and plant growth-promoting bacteria as phytoremediation methods increased plant tolerance at a metal-contaminated field site under low flood conditions.

546 **Behavior of *Trifolium repens* and *Lolium perenne* growing in a heavy metal contaminated field: Plant metal concentration and phytotoxicity**
G. Bidar, G. Garçon, C. Pruvot, D. Dewaele, F. Cazier, F. Douay, P. Shirali

Usefulness of Trifolium repens and Lolium perenne for the phytomanagement of heavy metal-contaminated soils.

554 **New exposure-based metric approach for evaluating O_3 risk to North American aspen forests**
K.E. Percy, M. Nosal, W. Heilman, T. Dann, J. Sober, A.H. Legge, D.F. Karnosky

A new exposure-based metric approach to predict O_3 risk to North American aspen forests has been developed.

567 **Drought changes the dynamics of trace element accumulation in a Mediterranean *Quercus ilex* forest**
J. Sardans, J. Peñuelas

Drought increased biomass concentrations of As and Cd and favors exportation of some trace elements to continental waters in a Mediterranean forest.

584 **Ca^{2+}-dependent plant response to Pb^{2+} is regulated by *LCT1***
S. Wojas, A. Ruszczyńska, E. Bulska, M. Wojciechowski, D.M. Antosiewicz

Ca^{2+}-dependent Pb^{2+} detoxification and uptake was regulated by TaLCT1.

593 **The leachability, bioaccessibility, and speciation of Cu in the sediment of channel catfish ponds**
R. Liu, D. Zhao

Careful disposal of the Cu-laden pond sediment is necessary.

604 **Toxicological responses of red-backed salamanders (*Plethodon cinereus*) to subchronic soil exposures of 2,4-dinitrotoluene**
M.S. Johnson, J. Suski, M.A. Bazar

Exposures of 2,4-dinitrotoluene in soil exceeding 345 mg/kg causes toxicity to P. cinereus.

609 **Dynamics of arbuscular mycorrhizal symbiosis in heavy metal phytoremediation: Meta-analytical and conceptual perspectives**
P. Audet, C. Charest

This meta-analysis has revealed a transition role of the AM symbiosis in phytoremediation shifting from 'Enhanced Uptake' to 'Metal-Binding' beyond critical soil-HM levels.

615 **Influence of hydrological regime on pore water metal concentrations in a contaminated sediment-derived soil**
G. Du Laing, D.R.J. Vanthuyne, B. Vandecasteele, F.M.G. Tack, M.G. Verloo

The hydrological regime is a key factor in determining the metal concentration in the pore water of a contaminated sediment-derived soil.

626 **Development and validation of a terrestrial biotic ligand model predicting the effect of cobalt on root growth of barley (*Hordeum vulgare*)**
K. Lock, K.A.C. De Schamphelaere, S. Becaus, P. Criel, H. Van Eeckhout, C.R. Janssen

Biotic Ligand Models are not only a useful tool to assess metal toxicity in aquatic systems but can also be used for terrestrial plants.

634 Halide salts accelerate degradation of high explosives by zerovalent iron
J.S. Kim, P.J. Shea, J.E. Yang, J.-E. Kim

The addition of halide ions promotes the degradation of high explosives by zerovalent iron.

642 Cadmium concentrations in tissues of willow ptarmigan (*Lagopus lagopus*) and rock ptarmigan (*Lagopus muta*) in Nunavik, Northern Québec
J. Rodrigue, L. Champoux, D. Leclair, J.-F. Duchesne

High levels of cadmium were found in ptarmigan in Northern Québec. No consumption guidelines are proposed for the Inuit people.

648 Evaluation of the treatment efficiencies of paper mill whitewaters in terms of organic composition and toxicity
A. Latorre, A. Malmqvist, S. Lacorte, T. Welander, D. Barceló

Aerobic and anaerobic treatments remove organic compounds in paper mill effluents but toxicity remains.

656 Linking valve closure behavior and sodium transport mechanism in freshwater clam *Corbicula fluminea* in response to copper
C.-M. Liao, C.-M. Lin, L.-J. Jou, K.-C. Chiang

A new flux–biological response model can link valve closure and sodium transport mechanisms in freshwater clam in response to copper.

668 Distribution and ecotoxicity of chlorotriazines in the Scheldt Estuary (B-NI)
H. Noppe, A. Ghekiere, T. Verslycke, E.D. Wulf, K. Verheyden, E. Monteyne, K. Polfliet, P. van Caeter, C.R. Janssen, H.F. De Brabander

The water of the Scheldt estuary and its associated suspended solids are contaminated with chlorotriazines at concentrations that do not affect mysid molting.

677 Tissue heavy metal concentrations of stranded California sea lions (*Zalophus californianus*) in Southern California
E.R. Harper, J.A. St. Leger, J.A. Westberg, L. Mazzaro, T. Schmitt, T.H. Reidarson, M. Tucker, D.H. Cross, B. Puschner

Tissue heavy metal concentrations are valuable in population and environmental monitoring.

683 Occurrence of alkylphenolic substances in a Great Lakes coastal marsh, Cootes Paradise, ON, Canada
T. Mayer, D. Bennie, F. Rosa, G. Rekas, V. Palabrica, J. Schachtschneider

Concentrations of alkylphenolic compounds in water, sediments and benthic invertebrates in a large coastal wetland and implications for trophic transfer.

691 *Psidium guajava* 'Paluma' (the guava plant) as a new bio-indicator of ozone in the tropics
C.M. Furlan, R.M. Moraes, P. Bulbovas, M. Domingos, A. Salatino, M.J. Sanz

Psidium guajava 'Paluma', a tropical species widely used in Brazilian food industry, is a potential sensitive bio-indicator of ozone.

696 Assessing airborne pollution effects on bryophytes – lessons learned through long-term integrated monitoring in Austria
H.G. Zechmeister, T. Dirnböck, K. Hülber, M. Mirtl

Bryophytes show ambiguous response to airborne pollutants during 14 years of monitoring in a forest ecosystem.

706 The use of sedimentary %C, %N, δ^{15}N, and Pb concentrations to assess historical changes in anthropogenic influence on Portuguese estuaries
P. Castro, I. Valiela, H. Freitas

The variables C, %N, δ^{15}N, and Pb were reliable sediment sentinels that reflected the level of anthropogenic influence in the past decades in estuarine ecosystems.

713 Estimating the extractability of potentially toxic metals in urban soils: A comparison of several extracting solutions
F. Madrid, R. Reinoso, M.C. Florido, E. Díaz Barrientos, F. Ajmone-Marsan, C.M. Davidson, L. Madrid

Dilute HCl extraction is tested as an alternative to the BCR sequential extraction in urban soils.

723 Root fungal colonisation in *Deschampsia flexuosa*: Effects of pollution and neighbouring trees
A.L. Ruotsalainen, A. Markkola, M.V. Kozlov

Severe pollution decreased root colonisation by some fungal groups; neighbouring trees decreased root colonisation by dark septate endophytic fungi in highly polluted sites.

729 Chromium and nickel mobilization from a contaminated soil using chelants
L. Jean, F. Bordas, J.-C. Bollinger

Chelant-induced dissolution of soil mineral matrix controls metal solubilization.

737 Effects of plant arsenic uptake and heavy metals on arsenic distribution in an arsenic-contaminated soil
A.O. Fayiga, L.Q. Ma, Q. Zhou

Arsenic taken up by P. vittata was from all fractions with most from the Ca-fraction.

743 Role of salicylic acid in alleviating oxidative damage in rice roots (*Oryza sativa*) subjected to cadmium stress
B. Guo, Y.C. Liang, Y.G. Zhu, F.J. Zhao

Pretreatment with SA enhanced the antioxidant defense activities in Cd-stressed rice, thus alleviating Cd-induced oxidative damage and enhancing Cd tolerance.

750 Effects of external phosphorus on the cell ultrastructure and the chlorophyll content of maize under cadmium and zinc stress
H.M. Jiang, J.C. Yang, J.F. Zhang

External P could decrease the bioavailability of Cd and Zn.

757 Validation of predicted exponential concentration profiles of chemicals in soils
A. Hollander, I. Baijens, A. Ragas, M. Huijbregts, D. van de Meent

Multimedia mass model predictions will improve when taking into account depth dependent soil concentrations.

764 Effect of lead on the sorption of 2,4,6-trichlorophenol on soil and peat
Z.-g. Pei, X.-q. Shan, T. Liu, Y.-n. Xie, B. Wen, S. Zhang, S.U. Khan

Lead diminished the sorption of 2,4,6-trichlorophenol onto soil and peat.

771 Multivariate statistical evaluation of trace elements in groundwater in a coastal area in Shenzhen, China
K. Chen, J.J. Jiao, J. Huang, R. Huang

Multivariate statistical analysis was used to investigate relationships among trace elements and factors controlling trace element distribution in groundwater.

781 Optimizing the molarity of a EDTA washing solution for saturated-soil remediation of trace metal contaminated soils
M.D. Andrade, S.O. Prasher, W.H. Hendershot

A single EDTA addition is best utilised in a highly concentrated washing solution given in a small liquid to soil weight ratio.

791 Use of ^{15}N-labelled nitrogen deposition to quantify the source of nitrogen in runoff at a coniferous-forested catchment at Gårdsjön, Sweden
O.J. Kjønaas, R.F. Wright

^{15}N tracer addition showed that initially the main source of NO_3^- in runoff was N from atmospheric deposition.

Short Communication

800 Physiological responses of *Alternanthera philoxeroides* (Mart.) Griseb leaves to cadmium stress
B. Ding, G. Shi, Y. Xu, J. Hu, Q. Xu

Cd modified a number of physiological processes and particularly chlorophyll degradation.

I Contents of Volume 147